

# List of Figures

2.1	Spinodal lines for the microphase transition and the macrophase separation in solutions of associating polymers . . . . .	20
2.2	Spinodal lines in the solution of associating polymers with chain length $N = 40$ and different end-block fractions, $f_A = 0.25, 0.3, 0.35$ . . . . .	21
2.3	Spinodal lines in the solution of associating polymers with large end blocks . . . . .	22
2.4	Critical wave vector in the solution of associating polymers . . . . .	23
2.5	Microphase spinodal calculated using the virial-type expansion instead of strict volume incompressibility . . . . .	24
3.1	Glass transitions in diblock-copolymer melt . . . . .	46
3.2	Chain-length dependence of glass transitions . . . . .	48
3.3	Glass transition vs ODT in symmetric copolymer melts . . . . .	51
3.4	Glass transition vs ODT in very asymmetric copolymer. (a) $\bar{N} = 10^7$ . (b) $\bar{N} = 10^8$ . .	52
3.5	Feynman diagrams: (a)–(d) Loop diagrams in $\Gamma_2$ . (e)–(g) Self-energy diagrams . . . . .	57
4.1	Schematic view of the model for surfaces with tethered ligands and receptors . . . . .	68
4.2	The contribution to the effective binding energy due to tether stretching . . . . .	84
4.3	The free energy of confining a polymer between parallel surfaces . . . . .	86
4.4	Comparison of case II (“both closed”) and case III (“open-closed”) . . . . .	88
4.5	Comparison of case III (“open-closed”) and case IV (“both open”) . . . . .	89
4.6	Comparison of case I (“quenched”) and case II (“both closed”) . . . . .	90
4.7	Dependence of the binding fraction and the free energy on the density of molecules for different cases . . . . .	92
4.8	Dependence of the equilibrium separation and free energy on the molecular binding energy for different cases . . . . .	93
4.9	The force-extension curve for different cases . . . . .	94
4.10	Schematic views of the models discussed in Section 4.3.3 . . . . .	97
4.11	The interaction between surfaces mediated by ligand-receptor binding and repulsive polymers . . . . .	99

4.12	The interaction potential resulting from binary ligand-receptor interactions . . . . .	101
4.13	Interaction between surfaces with two different ligand-receptor interactions and repelling polymers . . . . .	102
4.14	Green's function of joined rods . . . . .	108
5.1	Monge representation of a near-flat membrane shape . . . . .	121
5.2	Schematic representation of the interaction potential $V(z)$ . . . . .	123
5.3	Illustration of a regular adhesion droplet . . . . .	124
5.4	Shape of the potential $V(z)$ for $L_1 = L_2 = 1, V_1 = 2, V_2 = 1$ . . . . .	128
5.5	Evolution of the adhesion shape [(a) and (c)] and the free energy [(c) and (d)] along the minimum energy path for different adhesion potentials . . . . .	129
5.6	Critical membrane shapes at different potential depths . . . . .	131
5.7	The saddle point free energy $F^\ddagger$ and the critical shape radius $R^\ddagger$ at different potential depths . . . . .	132
5.8	The saddle point free energy $F^\ddagger$ and the critical shape radius $R^\ddagger$ at different potential well separations . . . . .	133
5.9	Illustration of the effects of surface tension . . . . .	135
5.10	Projection of an irregular droplet . . . . .	137

# List of Tables

4.1	Glossary . . . . .	70
4.2	Summary of different scenarios . . . . .	75
4.3	Scaled variables . . . . .	81
4.4	Examples of different cases . . . . .	87
5.1	Summary of scaling results . . . . .	139